

Influence™ LC

Fungicide

Registered
biofungicide for
powdery mildew
suppression on
Cannabis

Greenhouse Cannabis

FEATURES AND BENEFITS:

- ✓ **Zero** days to harvest, allows to keep control of diseases and maintain quality
- ✓ **Minimal** re-entry period (when spray is dry)
- ✓ **Safe** for use with beneficials and pollinators *
- ✓ **Minimum risk** of resistance
- ✓ **Perfect integration** in your IPM program

* Safe for use with beneficial organisms and pollinators when used according to label instructions.

**Also available in
smaller size starting
from 500 ml**



MODE OF ACTION

After treatment, conidia and mycelium of the pathogenic fungus growing on leaf surface are plasmolyzed and drying out.



AEF GLOBAL
Biopesticides

Influence™ LC

CROP	PEST	RATE	FOLIAR SPRAY VOLUME	APPLICATION TIMING
Cannabis (marihuana) produced commercially indoors	Powdery mildew	1.8 % (v/v) Max 170 ml/1000 pi²	Apply in sufficient water volume to ensure thorough leaves coverage. Use a minimum of 300 L/ha with a hydraulic sprayer.	Begin application at first sign of disease Repeat at 7-10 days interval under disease pressure

TECHNICAL SPECIFICATIONS

Active ingredient : Active ingredients: Garlic

Packaging: One (1) case = 2 x 10L, also available in smaller size starting from 500 ml

Formulation Type: Liquid concentrate

Chemical Group: N/A Biopesticide

PHI: 0 day

REI: When spray is dry



Influence LC is registered in Canada by AEF Global. Please refer to product label for complete application details. Always read and follow label directions. Log on to www.aefglobal.com for full label information.